

A new Lax pair for the sixth Painlevé equation associated with $\widehat{\mathfrak{so}}(8)$

Masatoshi NOUMI and Yasuhiko YAMADA

Department of Mathematics, Kobe University

Dedicated to Professor Mitsuo Morimoto on his sixtieth birthday

Introduction

In this article, we propose a new representation of Lax type for the sixth Painlevé equation. This representation, formulated in the framework of the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$ of type $D_4^{(1)}$, provides a natural explanation of the affine Weyl group symmetry of P_{VI} . After recalling a standard derivation of P_{VI} , we describe in Section 2 fundamental Bäcklund transformations for P_{VI} . In Section 3, we present our Lax pair for P_{VI} associated with $\mathfrak{so}(8)[z, z^{-1}]$, and explain how the Bäcklund transformations arise from the linear problem. For the general background on Painlevé equations, we refer the reader to [2].

The authors would like to thank Professor Kanehisa Takasaki for valuable discussions in the early stage of this work.

1 The sixth Painlevé equation

The sixth Painlevé equation is the following nonlinear ordinary differential equation of second order for the unknown function $y = y(t)$:

$$y'' = \frac{1}{2} \left(\frac{1}{y} + \frac{1}{y-1} + \frac{1}{y-t} \right) (y')^2 - \left(\frac{1}{t} + \frac{1}{t-1} + \frac{1}{y-t} \right) y' + \frac{y(y-1)(y-t)}{t^2(t-1)^2} \left(\alpha + \beta \frac{t}{y^2} + \gamma \frac{t-1}{(y-1)^2} + \delta \frac{t(t-1)}{(y-t)^2} \right), \quad (1)$$

where $'$ stands for the derivation with respect to the independent variable t , and $\alpha, \beta, \gamma, \delta$ are complex parameters.

A standard way to derive the sixth Painlevé equation is to employ the monodromy preserving deformation of a second order Fuchsian differential equation on $\mathbb{P}^1$, with four regular singular points and one apparent singularity. Consider the system of linear differential equations

$$(\partial_x^2 + a_1(x, t)\partial_x + a_2(x, t))u = 0, \quad \partial_t u = (b_1(x, t)\partial_x + b_2(x, t))u, \quad (2)$$

for the unknown functions $u = u(x, t)$, where $\partial_x = \partial/\partial x$ and $\partial_t = \partial/\partial t$. We assume that the coefficients $a_j(x, t)$ and $b_j(x, t)$ are rational functions in x , depending holomorphically on t , and that the first equation is Fuchsian with Riemann scheme

$$\left\{ \begin{array}{ccccc} x=0 & x=1 & x=t & x=q & x=\infty \\ 0 & 0 & 0 & 0 & \rho \\ \kappa_0 & \kappa_1 & \kappa_t & 2 & \kappa_\infty + \rho \end{array} \right\} \quad (3)$$

with respect to the variable x . In this scheme, $\kappa_0, \kappa_1, \kappa_t, \kappa_\infty$ and ρ are generic complex parameters subject to the *Fuchs relation*

$$\kappa_0 + \kappa_1 + \kappa_t + \kappa_\infty + 2\rho = 1. \quad (4)$$

We also assume that the singularity $x = q$, which may depend on t , is *non-logarithmic*. Under these assumptions, the coefficients $a_1(x, t)$, $a_2(x, t)$ are expressed in the form

$$\begin{aligned} a_1(x, t) &= \frac{1 - \kappa_0}{x} + \frac{1 - \kappa_1}{x - 1} + \frac{1 - \kappa_t}{x - t} - \frac{1}{x - q}, \\ a_2(x, t) &= \frac{1}{x(x - 1)} \left\{ -\frac{t(t - 1)H}{x - t} + \frac{q(q - 1)p}{x - q} + \rho(\kappa_\infty + \rho) \right\}, \end{aligned} \quad (5)$$

respectively, where

$$p = \text{Res}_{x=q}(a_2(x, t)dx), \quad H = -\text{Res}_{x=t}(a_2(x, t)dx). \quad (6)$$

Furthermore, the coefficient H is determined as a polynomial in (q, p) with coefficients in $\mathbb{C}(t)$; explicitly, it is given by

$$\begin{aligned} H &= \frac{1}{t(t - 1)} \left[p^2 q(q - 1)(q - t) - p \{ \kappa_0(q - 1)(q - t) \right. \\ &\quad \left. + \kappa_1 q(q - t) + (\kappa_t - 1)q(q - 1) \} + \rho(\kappa_\infty + \rho)(q - t) \right]. \end{aligned} \quad (7)$$

The compatibility condition of the linear differential system (2) then turns out to be expressed as the Hamiltonian system

$$H_{\text{VI}} : \quad \partial_t(q) = \frac{\partial H}{\partial p}, \quad \partial_t(p) = -\frac{\partial H}{\partial q}, \quad (8)$$

with polynomial Hamiltonian H in (7); namely,

$$\begin{aligned} t(t - 1)\partial_t(q) &= 2pq(q - 1)(q - t) - \{ \kappa_0(q - 1)(q - t) \\ &\quad + \kappa_1 q(q - t) + (\kappa_t - 1)q(q - 1) \} \\ t(t - 1)\partial_t(p) &= -p^2(3q^2 - 2(1 + t)q + t) + p \{ 2(\kappa_0 + \kappa_1 + \kappa_t - 1)q \\ &\quad - \kappa_0(1 + t) - \kappa_1 t - \kappa_t + 1 \} - \rho(\kappa_\infty + \rho). \end{aligned} \quad (9)$$

This system of nonlinear equations is in fact equivalent to the sixth Painlevé equation P_{VI} (1) for $y = q$, with parameters

$$\alpha = \frac{\kappa_\infty^2}{2}, \quad \beta = -\frac{\kappa_0^2}{2}, \quad \gamma = \frac{\kappa_1^2}{2}, \quad \delta = \frac{1 - \kappa_t^2}{2}. \quad (10)$$

We remark that, in place of (2), one can naturally make use of the Schlesinger system of rank two, with regular singular points $x = 0, 1, t, \infty$.

2 Discrete symmetry of H_{VI}

It is known that the sixth Painlevé equation admits a group of Bäcklund transformations which is isomorphic to the (extended) affine Weyl group of type $D_4^{(1)}$ (see [7], for instance).

In describing the Bäcklund transformations for H_{VI} , it is convenient to use the parameters

$$\alpha_0 = \kappa_t, \quad \alpha_1 = \kappa_\infty, \quad \alpha_2 = \rho, \quad \alpha_3 = \kappa_1, \quad \alpha_4 = \kappa_0 \quad (11)$$

with linear relation $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$, so that

$$\begin{aligned} t(t-1)H = & p^2q(q-1)(q-t) - p\{(\alpha_0-1)q(q-1) \\ & + \alpha_3q(q-t) + \alpha_4(q-1)(q-t)\} + \alpha_2(\alpha_1 + \alpha_2)(q-t). \end{aligned} \quad (12)$$

In the following, we identify the parameter space for H_{VI} with the affine space $V = \mathbb{C}^4$ with canonical coordinates $\varepsilon = (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4)$, and regard α_j as linear functions of V such that

$$\begin{aligned} \alpha_0 &= 1 - \varepsilon_1 - \varepsilon_2, & \alpha_1 &= \varepsilon_1 - \varepsilon_2, & \alpha_2 &= \varepsilon_2 - \varepsilon_3, \\ \alpha_3 &= \varepsilon_3 - \varepsilon_4, & \alpha_4 &= \varepsilon_3 + \varepsilon_4. \end{aligned} \quad (13)$$

We identify V with the Cartan subalgebra of the simple Lie algebra $\mathfrak{so}(8)$ (of type D_4); $\{\varepsilon_1, \dots, \varepsilon_4\}$ is then a canonical orthonormal basis of V^* , and $\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4$ are the *simple affine roots*. Note that the null root $\delta = \alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4$ is normalized to be the constant function 1.

By a *Bäcklund transformation*, we mean a transformation of dependent variables and parameters that leaves the system invariant. Let us show an example of Bäcklund transformation for H_{VI} . Define new variables $\tilde{q}, \tilde{p}$ by

$$\tilde{q} = q, \quad \tilde{p} = p - \frac{\alpha_0}{q-t}. \quad (14)$$

Then one can verify directly that, if the pair (q, p) satisfies the Hamiltonian system (8), then the pair $(\tilde{q}, \tilde{p})$ again satisfies the same system with parameters α_0, α_2 replaced by $-\alpha_0, \alpha_2 + \alpha_0$, respectively; we refer to this Bäcklund transformation as s_0 . To be more precise, let us consider the field of rational functions

$$\mathcal{K} = \mathbb{C}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, q, p, t) \quad (\alpha_0 = 1 - \alpha_1 - 2\alpha_2 - \alpha_3 - \alpha_4), \quad (15)$$

and the Hamiltonian vector field

$$\delta = \frac{\partial H}{\partial p} \frac{\partial}{\partial q} - \frac{\partial H}{\partial q} \frac{\partial}{\partial p} + \frac{\partial}{\partial t} \quad (16)$$

acting on $\mathcal{K}$ as a derivation. We regard this *differential field* $(\mathcal{K}, \delta)$ as representing the Hamiltonian system H_{VI} . We define the automorphism $s_0 : \mathcal{K} \rightarrow \mathcal{K}$ by setting

$$s_0(\alpha_0) = -\alpha_0, \quad s_0(\alpha_2) = \alpha_2 + \alpha_0, \quad s_0(\alpha_j) = \alpha_j \quad (j \neq 0, 2), \quad (17)$$

Table 1: Bäcklund transformations for H_{VI}

	α_0	α_1	α_2	α_3	α_4	q	p
s_0	$-\alpha_0$	α_1	$\alpha_2 + \alpha_0$	α_3	α_4	q	$p - \frac{\alpha_0}{q-t}$
s_1	α_0	$-\alpha_1$	$\alpha_2 + \alpha_1$	α_3	α_4	q	p
s_2	$\alpha_0 + \alpha_2$	$\alpha_1 + \alpha_2$	$-\alpha_2$	$\alpha_3 + \alpha_2$	$\alpha_4 + \alpha_2$	$q + \frac{\alpha_2}{p}$	p
s_3	α_0	α_1	$\alpha_2 + \alpha_3$	$-\alpha_3$	α_4	q	$p - \frac{\alpha_3}{q-1}$
s_4	α_0	α_1	$\alpha_2 + \alpha_4$	α_3	$-\alpha_4$	q	$p - \frac{\alpha_4}{q}$
r_1	α_1	α_0	α_2	α_4	α_3	$\frac{t(q-1)}{q-t}$	$-\frac{(q-t)((q-t)p+\alpha_2)}{t(t-1)}$
r_3	α_3	α_4	α_2	α_0	α_1	$\frac{t}{q}$	$-\frac{q(qp+\alpha_2)}{t}$
r_4	α_4	α_3	α_2	α_1	α_3	$\frac{q-t}{q-1}$	$\frac{(q-1)((q-1)p+\alpha_2)}{t-1}$

and

$$s_0(q) = q, \quad s_0(p) = p - \frac{\alpha_0}{q-t}, \quad s_0(t) = t. \quad (18)$$

Then one can show that the automorphism $s_0 : \mathcal{K} \rightarrow \mathcal{K}$ commutes with the Hamiltonian vector field δ . In this sense, a Bäcklund transformation can be defined alternatively to be an automorphism of the differential field that commutes with the derivation.

Table 1 is the list of *fundamental* Bäcklund transformations for H_{VI} . We consider two subgroups

$$W = \langle s_0, s_1, s_2, s_3, s_4 \rangle \subset \widetilde{W} = \langle s_0, s_1, s_2, s_3, s_4, r_1, r_3, r_4 \rangle \subset \text{Aut}_\delta(\mathcal{K}) \quad (19)$$

of differential automorphisms of $\mathcal{K}$, generated by the Bäcklund transformations in Table 1. Then it turns out that W and $\widetilde{W}$ are isomorphic to the affine Weyl group and the extended affine Weyl group of type $D_4^{(1)}$, respectively. The Bäcklund transformations s_i ($i = 0, 1, 2, 3, 4$) and r_i ($i = 1, 3, 4$) in fact satisfy the fundamental relations

$$\begin{aligned}
 s_i^2 &= 1 & (i = 0, 1, 2, 3, 4), \\
 s_i s_j &= s_j s_i & (i, j = 0, 1, 3, 4), \\
 s_i s_2 s_i &= s_2 s_i s_2 & (i = 0, 1, 3, 4), \\
 r_i^2 &= 1 & (i = 1, 3, 4) \\
 r_i r_j &= r_k & (\{i, j, k\} = \{1, 3, 4\}) \\
 r_i s_j &= s_{\sigma_i(j)} r_i & (i = 1, 3, 4; j = 0, 1, 2, 3, 4),
 \end{aligned} \quad (20)$$

where σ_i ($i = 1, 3, 4$) are the permutations defined by

$$\sigma_1 = (01)(34), \quad \sigma_3 = (03)(14), \quad \sigma_4 = (04)(13). \quad (21)$$

We also remark that each element $w \in \widetilde{W}$ defines a *canonical transformation*:

$$w(\{\varphi, \psi\}) = \{w(\varphi), w(\psi)\} \quad (\varphi, \psi \in \mathcal{K}), \quad (22)$$

where $\{, \}$ stands for the standard Poisson bracket defined by

$$\{\varphi, \psi\} = \frac{\partial \varphi}{\partial p} \frac{\partial \psi}{\partial q} - \frac{\partial \varphi}{\partial q} \frac{\partial \psi}{\partial p}. \quad (23)$$

Remark 2.1 The fundamental relations for the generators s_0, s_1, s_2, s_3, s_4 of the affine Weyl group of type $D_4^{(1)}$ is described as follows in terms of the Cartan matrix $A = (a_{ij})_{i,j=0}^4$:

$$\begin{aligned} s_i^2 &= 1 & (i = 0, 1, 2, 3, 4) \\ s_i s_j &= s_j s_i & \text{if } (a_{ij}, a_{ji}) = (0, 0), \\ s_i s_j s_i &= s_j s_i s_j & \text{if } (a_{ij}, a_{ji}) = (-1, -1), \end{aligned} \quad (24)$$

where

$$A = \begin{bmatrix} 2 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 \\ -1 & -1 & 2 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 2 \end{bmatrix}. \quad \begin{array}{c} \circ 0 \quad \circ 3 \\ \diagdown \quad \diagup \\ \circ 2 \\ \diagup \quad \diagdown \\ \circ 1 \quad \circ 4 \end{array} \quad (25)$$

The action of s_i on the simple affine roots α_j is given by

$$s_i(\alpha_j) = \alpha_j - \alpha_i a_{ij} \quad (i, j = 0, 1, 2, 3, 4). \quad (26)$$

Note also that $\widetilde{W}$ is isomorphic to the semidirect product $W \rtimes \Omega$ of W and $\Omega = \{1, r_1, r_3, r_4\}$ acting on W through the permutations $\{1, \sigma_1, \sigma_3, \sigma_4\}$ of indices for the generators s_j ; Ω is identified with a group of diagram automorphisms of the Dynkin diagram of type $D_4^{(1)}$. If we set

$$\varphi_0 = q - t, \quad \varphi_1 = 1, \quad \varphi_2 = -p, \quad \varphi_3 = q - 1, \quad \varphi_4 = q, \quad (27)$$

the Bäcklund transformations s_i are expressed as

$$s_i(\varphi_j) = \varphi_j + \frac{\alpha_i}{\varphi_i} u_{ij}, \quad u_{ij} = \{\varphi_i, \varphi_j\} \quad (i, j = 0, 1, 2, 3, 4), \quad (28)$$

consistently with the birational Weyl group actions discussed previously in [5], [6]. In this particular case, the matrix $U = (u_{ij})_{i,j=0}^4$ is given by

$$U = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix}. \quad (29)$$

Observe that $u_{12} = u_{21} = 0$; this degeneracy is caused by the normalization that one of the regular singular points of (2) is placed at $x = \infty$.

Remark 2.2 The extended affine Weyl group $\widetilde{W}$ of type $D_4^{(1)}$ is expressed as the semidirect product of the weight lattice P of type D_4 and the Weyl group $W(D_4) = \langle s_1, s_2, s_3, s_4 \rangle$ acting on P :

$$\widetilde{W} \simeq P \rtimes W(D_4), \quad P = \bigoplus_{i=1}^4 \mathbb{Z} \varpi_i, \quad (30)$$

where ϖ_i are the fundamental weights of type D_4 defined by

$$\begin{aligned} \varpi_1 &= \varepsilon_1, & \varpi_2 &= \varepsilon_1 + \varepsilon_2, \\ \varpi_3 &= \frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 - \varepsilon_4), & \varpi_4 &= \frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4). \end{aligned} \quad (31)$$

Note also that the weight lattice P is the set of all elements

$$\lambda = \frac{1}{2}(n_1\varepsilon_1 + n_2\varepsilon_2 + n_3\varepsilon_3 + n_4\varepsilon_4) \quad (n_1, n_2, n_3, n_4 \in \mathbb{Z}) \quad (32)$$

such that, either all the n_j 's are even, or all the n_j 's are odd. The translations T_{ϖ_i} ($i = 1, \dots, 4$) corresponding to ϖ_i are expressed as

$$\begin{aligned} T_{\varpi_1} &= r_1 s_1 s_2 s_3 s_4 s_2 s_1, & T_{\varpi_2} &= s_0 s_2 s_1 s_3 s_4 s_2 s_1 s_3 s_4 s_2, \\ T_{\varpi_3} &= r_3 s_3 s_2 s_1 s_4 s_2 s_3, & T_{\varpi_4} &= r_4 s_4 s_2 s_1 s_3 s_2 s_4, \end{aligned} \quad (33)$$

in terms of the generators s_j and r_j . These elements transform the simple affine roots α_j as follows.

	α_0	α_1	α_2	α_3	α_4
T_{ϖ_1}	$\alpha_0 + 1$	$\alpha_1 - 1$	α_2	α_3	α_4
T_{ϖ_2}	$\alpha_0 + 2$	α_1	$\alpha_2 - 1$	α_3	α_4
T_{ϖ_3}	$\alpha_0 + 1$	α_1	α_2	$\alpha_3 - 1$	α_4
T_{ϖ_4}	$\alpha_0 + 1$	α_1	α_2	α_3	$\alpha_4 - 1$

(34)

Regarded as automorphisms of $\mathcal{K} = \mathbb{C}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, q, p, t)$, T_{ϖ_i} ($i = 1, 2, 3, 4$) provide a commuting family of Bäcklund transformations for H_{VI} , which are called *Schlesinger transformations* (such Bäcklund transformations that act as shift operators on the parameter space).

A certain part of this discrete symmetry of H_{VI} can be explained by the monodromy preserving deformation of a second order Fuchsian equation (2). In fact, each s_i ($i = 0, 1, 3, 4$), except s_2 , arises from a simple transformation of the unknown function $u = u(x, t)$, and each r_i ($i = 1, 3, 4$) from a fractional linear transformation of the coordinate x . This framework does not seem, however, to explain the particular Bäcklund transformation s_2 in Table 1, which is essential in understanding the whole picture of discrete symmetry of H_{VI} . In the next section, we propose a new Lax pair for H_{VI} , from which all the Bäcklund transformations in Table 1 can be understood naturally.

3 Lax pair associated with $\widehat{\mathfrak{so}}(8)$

Consider the following system of linear differential equations for the column vector $\vec{\psi} = (\psi_1, \psi_2, \dots, \psi_8)^t$ of eight unknown functions $\psi_i = \psi_i(z, t)$ ($i = 1, 2, \dots, 8$):

$$(z\partial_z + M)\vec{\psi} = 0, \quad \partial_t \vec{\psi} = B\vec{\psi}, \quad (35)$$

with the compatibility condition

$$[z\partial_z + M, \partial_t - B] = 0. \quad (36)$$

We assume that the matrices M and B are in the form

$$M = \begin{bmatrix} \varepsilon_1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \varepsilon_2 & -p & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & \varepsilon_3 & q-1 & q & 0 & 0 & 0 \\ 0 & 0 & 0 & \varepsilon_4 & 0 & -q & 1 & 0 \\ 0 & 0 & 0 & 0 & -\varepsilon_4 & 1-q & 1 & 0 \\ -z & 0 & 0 & 0 & 0 & -\varepsilon_3 & p & 0 \\ (t-q)z & 0 & 0 & 0 & 0 & 0 & -\varepsilon_2 & -1 \\ 0 & (q-t)z & z & 0 & 0 & 0 & 0 & -\varepsilon_1 \end{bmatrix} \quad (37)$$

and

$$B = \begin{bmatrix} u_1 & x_1 & y_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & u_2 & x_2 & -y_3 & -y_4 & 0 & 0 & 0 \\ 0 & 0 & u_3 & x_3 & x_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & u_4 & 0 & -x_4 & y_4 & 0 \\ 0 & 0 & 0 & 0 & -u_4 & -x_3 & y_3 & 0 \\ 0 & 0 & 0 & 0 & 0 & -u_3 & -x_2 & -y_1 \\ -z & 0 & 0 & 0 & 0 & 0 & -u_2 & -x_1 \\ 0 & z & 0 & 0 & 0 & 0 & 0 & -u_1 \end{bmatrix}, \quad (38)$$

respectively, where ε_j are complex constants, and the variables q, p, x_j, y_j and u_j are functions in t . As before, we set

$$\begin{aligned} \alpha_0 &= 1 - \varepsilon_1 - \varepsilon_2, & \alpha_1 &= \varepsilon_1 - \varepsilon_2, & \alpha_2 &= \varepsilon_2 - \varepsilon_3, \\ \alpha_3 &= \varepsilon_3 - \varepsilon_4, & \alpha_4 &= \varepsilon_3 + \varepsilon_4. \end{aligned} \quad (39)$$

Theorem 3.1 *Under the compatibility condition (36), the variables x_j , y_j and u_j are determined uniquely as elements of $\mathcal{K} = \mathbb{C}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, q, p, t)$. The compatibility condition is then equivalent to the Hamiltonian system H_{V1} of the sixth Painlevé equation.*

The variables x_j , y_j and u_j appearing in B are determined explicitly as follows:

$$\begin{aligned} x_1 &= \frac{q-t}{t(t-1)}, & x_3 &= -\frac{q}{t}, & x_4 &= -\frac{q-1}{t-1}, \\ y_1 &= \frac{1}{t(t-1)}, & y_3 &= -\frac{1}{t}, & y_4 &= -\frac{1}{t-1}, \\ x_2 &= -\frac{(q-t)p + \alpha_1 + \alpha_2}{t(t-1)}, \end{aligned} \quad (40)$$

and

$$\begin{aligned}
t(t-1)u_1 &= -q(q-1)p - \alpha_2 q + \frac{\alpha_0 - \alpha_1 - 1}{2}t - \frac{\alpha_0 + \alpha_4 - 1}{2}, \\
t(t-1)u_2 &= -q(q-1)p - (\alpha_1 + \alpha_2)q + \frac{\alpha_0 + \alpha_1 - 1}{2}t - \frac{\alpha_0 + \alpha_4 - 1}{2}, \\
t(t-1)u_3 &= (2q-1)(q-t)p + (\alpha_1 + 2\alpha_2)q + \frac{\alpha_3 + \alpha_4}{2}t + \frac{\alpha_0 + \alpha_4 - 1}{2}, \\
t(t-1)u_4 &= -(q-t)p + \frac{\alpha_3 - \alpha_4}{2}t + \frac{\alpha_0 + \alpha_4 - 1}{2}.
\end{aligned} \tag{41}$$

We now explain how this representation is related to the Lie algebra $\mathfrak{so}(8)$ and its loop algebra. With the notation of matrix units $E_{ij} = (\delta_{ia}\delta_{jb})_{a,b=1}^8$, we set

$$J = \sum_{i=1}^8 E_{i,9-i}. \tag{42}$$

We consider the following realization of the simple Lie algebra $\mathfrak{so}(8)$:

$$\mathfrak{so}(8) = \{X \in \text{Mat}(8; \mathbb{C}) \mid JX + X^t J = 0\}, \tag{43}$$

where X^t denotes the transposition of X . Let us define the Chevalley generators E_j, H_j, F_j ($j = 0, 1, 2, 3, 4$) for the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$ by

$$\begin{aligned}
E_0 &= z(E_{82} - E_{71}), & E_1 &= E_{12} - E_{78}, & E_2 &= E_{23} - E_{67}, \\
E_3 &= E_{34} - E_{56}, & E_4 &= E_{35} - E_{46}, \\
F_0 &= z^{-1}(E_{28} - E_{17}), & F_1 &= E_{21} - E_{87}, & F_2 &= E_{32} - E_{76}, \\
F_3 &= E_{43} - E_{65}, & F_4 &= E_{53} - E_{64},
\end{aligned} \tag{44}$$

and $H_j = [E_j, F_j]$ ($j = 0, 1, 2, 3, 4$). For a vector $\mathbf{a} = (a_1, a_2, a_3, a_4)$ given, we also use the notation

$$H(\mathbf{a}) = \sum_{i=1}^4 a_i (E_{ii} - E_{9-i,9-i}) \tag{45}$$

for the corresponding element in the Cartan subalgebra of $\mathfrak{so}(8)$, so that

$$\begin{aligned}
H_0 &= H(-1, -1, 0, 0), & H_1 &= H(1, -1, 0, 0), & H_2 &= H(0, 1, -1, 0) \\
H_3 &= H(0, 0, 1, -1), & H_4 &= H(0, 0, 1, 1).
\end{aligned} \tag{46}$$

Notice that the two matrices M, B belong to a Borel subalgebra of the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$; in fact, they are expressed in the form

$$\begin{aligned}
M &= H(\boldsymbol{\varepsilon}) + (q-t)E_0 + E_1 - pE_2 + (q-1)E_3 + qE_4 \\
&\quad + [E_0, E_2] + [E_3, E_2] + [E_4, E_2], \\
B &= H(\mathbf{u}) + E_0 + x_1E_1 + x_2E_2 + x_3E_3 + x_4E_4 \\
&\quad + y_1[E_1, E_2] + y_3[E_3, E_2] + y_4[E_4, E_2],
\end{aligned} \tag{47}$$

where $\boldsymbol{\varepsilon} = (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4)$ and $\mathbf{u} = (u_1, u_2, u_3, u_4)$.

Remark 3.2 The affine Lie algebra $\widehat{\mathfrak{so}}(8)$ is realized as a central extension of the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$, together with the derivation $d = z\partial_z$ (see [4], for the detail) :

$$\widehat{\mathfrak{so}}(8) = \mathfrak{so}(8) \otimes \mathbb{C}[z, z^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d, \quad (48)$$

where c denotes the canonical central element. In this context, the simple affine roots α_j ($j = 0, 1, 2, 3, 4$) are defined as linear functionals of the Cartan subalgebra

$$\mathfrak{h} = \bigoplus_{i=0}^4 \mathbb{C}h_i \oplus \mathbb{C}d, \quad h_0 = H_0 \otimes 1 + c, \quad h_i = H_i \otimes 1 \quad (i = 1, 2, 3, 4) \quad (49)$$

such that

$$\langle h_i, \alpha_j \rangle = a_{i,j} \quad (i = 0, 1, 2, 3, 4), \quad \langle d, \alpha_j \rangle = \delta_{0,j} \quad (50)$$

for $j = 0, 1, 2, 3, 4$; also they are extended to linear functionals on the whole affine Lie algebra $\widehat{\mathfrak{so}}(8)$ through the triangular decomposition. Note that our Lax pair mentioned above is formulated in fact in the framework of $\widehat{\mathfrak{so}}(8)/\mathbb{C}c = \mathfrak{so}(8)[z, z^{-1}] \oplus \mathbb{C}d$:

$$\mathcal{M} = d + M = z\partial_z + M \in \widehat{\mathfrak{so}}(8)/\mathbb{C}c. \quad (51)$$

Since $\langle c, \alpha_j \rangle = 0$, we can regard α_j ($j = 0, 1, 2, 3, 4$) and the null root $\delta = \alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4$ as linear functionals on $\widehat{\mathfrak{so}}(8)/\mathbb{C}c$. Then we have

$$\begin{aligned} \alpha_0(\mathcal{M}) &= 1 - \varepsilon_1 - \varepsilon_2, & \alpha_1(\mathcal{M}) &= \varepsilon_1 - \varepsilon_2, & \alpha_2(\mathcal{M}) &= \varepsilon_2 - \varepsilon_3, \\ \alpha_3(\mathcal{M}) &= \varepsilon_3 - \varepsilon_4, & \alpha_4(\mathcal{M}) &= \varepsilon_3 + \varepsilon_4, & \delta(\mathcal{M}) &= 1. \end{aligned} \quad (52)$$

In this sense, our notation for the parameters (39) is consistent with that of simple roots for $\widehat{\mathfrak{so}}(8)$.

In our framework, the Bäcklund transformations for H_{VI} are obtained as the gauge transformations

$$s_k \vec{\psi} = G_k \vec{\psi} \quad (k = 0, 1, 2, 3, 4), \quad r_k \vec{\psi} = \Gamma_k \vec{\psi} \quad (k = 1, 3, 4) \quad (53)$$

of the linear problem (35), defined by certain matrices G_k, Γ_k in the loop group of

$$SO(8) = \{X \in SL(8; \mathbb{C}) \mid X^t J X = J\}. \quad (54)$$

The matrices G_k and Γ_k will be specified below.

Theorem 3.3 *The Bäcklund transformations s_k and r_k for H_{VI} are recovered from the compatibility conditions*

$$s_k(M) = G_k M G_k^{-1} - z\partial_z(G_k)G_k^{-1}, \quad s_k(B) = G_k B G_k^{-1} + \partial_t(G_k)G_k^{-1}, \quad (55)$$

and

$$r_k(M) = \Gamma_k M \Gamma_k^{-1} - z\partial_z(\Gamma_k)\Gamma_k^{-1}, \quad r_k(B) = \Gamma_k B \Gamma_k^{-1} + \partial_t(\Gamma_k)\Gamma_k^{-1}. \quad (56)$$

In (53), the matrices G_k are determined as

$$\begin{aligned} G_0 &= 1 + \frac{\alpha_0}{q-t} F_0, & G_1 &= 1 + \alpha_1 F_1, & G_2 &= 1 - \frac{\alpha_2}{p} F_2, \\ G_3 &= 1 + \frac{\alpha_3}{q-1} F_3, & G_4 &= 1 + \frac{\alpha_4}{q} F_4. \end{aligned} \quad (57)$$

The matrices Γ_1, Γ_3 and Γ_4 are given explicitly by

$$\Gamma_1 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{z\sqrt{t(t-1)}} \\ 0 & \frac{q-t}{\sqrt{t(t-1)}} & \frac{1}{\sqrt{t(t-1)}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{\sqrt{t(t-1)}}{q-t} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\frac{t}{\sqrt{t(t-1)}} & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{\sqrt{t(t-1)}}{t} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\frac{q-t}{\sqrt{t(t-1)}} & \frac{1}{\sqrt{t(t-1)}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{t(t-1)}}{q-t} & 0 \\ z\sqrt{t(t-1)} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad (58)$$

$$\Gamma_3 = \begin{bmatrix} 0 & 0 & 0 & \frac{1}{\sqrt{-t}\sqrt{z}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{-q}{\sqrt{-t}\sqrt{z}} & \frac{1}{\sqrt{-t}\sqrt{z}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{-t}}{q\sqrt{z}} & 0 \\ \sqrt{-t}\sqrt{z} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{-t}\sqrt{z}} \\ 0 & \frac{q\sqrt{z}}{\sqrt{-t}} & \frac{\sqrt{z}}{\sqrt{-t}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\frac{\sqrt{-t}\sqrt{z}}{q} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{-t}\sqrt{z} & 0 & 0 & 0 \end{bmatrix}, \quad (59)$$

$$\Gamma_4 = \begin{bmatrix} 0 & 0 & 0 & 0 & \frac{1}{\sqrt{1-t}\sqrt{z}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1-q}{\sqrt{1-t}\sqrt{z}} & \frac{1}{\sqrt{1-t}\sqrt{z}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{1-t}}{(1-q)\sqrt{z}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{\sqrt{1-t}\sqrt{z}} \\ \sqrt{1-t}\sqrt{z} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{(1-q)\sqrt{z}}{\sqrt{1-t}} & -\frac{\sqrt{z}}{\sqrt{1-t}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{\sqrt{1-t}\sqrt{z}}{1-q} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sqrt{1-t}\sqrt{z} & 0 & 0 & 0 & 0 \end{bmatrix}. \quad (60)$$

We remark that the matrices Γ_k ($k = 1, 3, 4$) are expressed as

$$\begin{aligned}\Gamma_1 &= D(\mathbf{a}_1) \exp\left(\frac{1}{q-t} E_2\right) z^{-\varpi_1} C_1, \\ \Gamma_3 &= D(\mathbf{a}_2) \exp\left(\frac{1}{q} E_2\right) z^{-\varpi_3} C_3, \\ \Gamma_4 &= D(\mathbf{a}_3) \exp\left(\frac{1}{q-1} E_2\right) z^{-\varpi_4} C_4,\end{aligned}\tag{61}$$

where we have used the notation

$$D(\mathbf{a}) = \text{diag}(a_1, a_2, a_3, a_4, a_4^{-1}, a_3^{-1}, a_2^{-1}, a_1^{-1})\tag{62}$$

for a vector $\mathbf{a} = (a_1, a_2, a_3, a_4)$, $a_j \neq 0$. In (61), z^{ϖ_k} denote the following diagonal matrices associated with the fundamental weights of $\mathfrak{so}(8)$:

$$\begin{aligned}z^{\varpi_1} &= z^{H(1,0,0,0)} = D(z, 1, 1, 1), \\ z^{\varpi_2} &= z^{H(1,1,0,0)} = D(z, z, 1, 1), \\ z^{\varpi_3} &= z^{H(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2})} = D(z^{\frac{1}{2}}, z^{\frac{1}{2}}, z^{\frac{1}{2}}, z^{-\frac{1}{2}}), \\ z^{\varpi_4} &= z^{H(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})} = D(z^{\frac{1}{2}}, z^{\frac{1}{2}}, z^{\frac{1}{2}}, z^{\frac{1}{2}}).\end{aligned}\tag{63}$$

The matrices C_k are essentially permutation matrices; with the notation of permutation matrices $S_\sigma = (\delta_{\sigma(i),j})_{i,j=1}^8$ for $\sigma \in \mathbf{S}_8$,

$$\begin{aligned}C_1 &= S_{(18)(45)}, \\ C_3 &= D(1, -1, 1, -1) S_{(14)(26)(37)(58)}, \\ C_4 &= D(1, -1, 1, -1) S_{(15)(26)(37)(48)}.\end{aligned}\tag{64}$$

The matrices $D(\mathbf{a}_1), D(\mathbf{a}_3), D(\mathbf{a}_4)$ in (61) are defined by

$$\begin{aligned}\mathbf{a}_1 &= \left(\frac{1}{\sqrt{t(t-1)}}, \frac{q-t}{\sqrt{t(t-1)}}, -\frac{\sqrt{t(t-1)}}{q-t}, \frac{-t}{\sqrt{t(t-1)}} \right), \\ \mathbf{a}_3 &= \left(\frac{1}{\sqrt{-t}}, \frac{q}{\sqrt{-t}}, \frac{\sqrt{-t}}{q}, -\sqrt{-t} \right), \\ \mathbf{a}_4 &= \left(\frac{1}{\sqrt{1-t}}, \frac{q-1}{\sqrt{1-t}}, \frac{\sqrt{1-t}}{1-q}, \frac{-1}{\sqrt{1-t}} \right).\end{aligned}\tag{65}$$

We also remark that, for each $k = 1, 3, 4$, the adjoint action of the matrix $z^{-\varpi_k} C_k$ induces the automorphism of the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$ corresponding to the diagram automorphism $\sigma_1 = (01)(34)$, $\sigma_3 = (03)(14)$ or $\sigma_4 = (04)(13)$, respectively. The remaining part of Γ_k concerns the normalization of the matrices M and B .

Note that the system of differential equations (35) has a regular singularity at $z = 0$ with exponents $\pm \varepsilon_j$ ($j = 1, 2, 3, 4$), and an irregular singularity at $z = \infty$. Assuming that ε_j are generic, let us take a fundamental system of solutions $\Psi = \Psi(z, t)$ of (35) with normalization such that

$$\Psi(z, t) = \sum_{n=0}^{\infty} \Psi_n(t) z^{-H(\varepsilon)+n}\tag{66}$$

around $z = 0$, and that $\Psi_0(t)$ is upper triangular; such a Ψ is determined up to the multiplication of constant diagonal matrices. Then, for each $k = 0, 1, 2, 3, 4$, the Bäcklund transformation s_k is interpreted as the transformation

$$\Psi \mapsto \tilde{\Psi} = G_k \Psi S_k \quad (67)$$

of the fundamental system of solutions, where $S_k = \exp(-E_k) \exp(F_k) \exp(-E_k)$ denote a lift of s_k to the loop group of $SO(8)$. Similarly, for each $k = 1, 3, 4$, the Bäcklund transformation r_k is interpreted as the transformation

$$\Psi \mapsto \tilde{\Psi} = \Gamma_k \Psi R_k, \quad (68)$$

where $R_k = z^{-\varpi_k} C_k$.

Remark 3.4 The system of differential equations (35) can be equivalently rewritten into a chain of systems of rank 2. We first extend the indexing set for ψ_i and ε_i to $\mathbb{Z}$ by imposing the periodicity condition

$$\psi_{i+8} = z\psi_i, \quad \varepsilon_{i+8} = \varepsilon_i - 1, \quad \varepsilon_{9-i} = -\varepsilon_i \quad (i \in \mathbb{Z}). \quad (69)$$

Then (35) is equivalent to a system for the 2-vectors $\vec{\psi}_i = (\psi_{2i}, \psi_{2i+1})^t$ ($i \in \mathbb{Z}$) in the following form:

$$(z\partial_z + M_i)\vec{\psi}_i + N_i\vec{\psi}_{i+1} = 0, \quad (-\partial_t + A_i)\vec{\psi}_i + B_i\vec{\psi}_{i+1} = 0, \quad (70)$$

where M_i, N_i, A_i, B_i are 2×2 matrices whose entries depend only on t . This system is formally transformed into

$$(\lambda + M_i)\vec{\varphi}_i + N_i\vec{\varphi}_{i+1} = 0, \quad (-\partial_t + A_i)\vec{\varphi}_i + B_i\vec{\varphi}_{i+1} = 0, \quad (71)$$

where $\vec{\varphi}_i = \vec{\varphi}_i(\lambda, t)$, by the change of coordinates $z = e^w$ and the Laplace transformation $\partial_w \leftrightarrow \lambda$, $w \leftrightarrow -\partial_\lambda$. Noting that N_i are invertible, one can rewrite (71) as

$$\partial_t \vec{\varphi}_i = U_i \vec{\varphi}_i, \quad \vec{\varphi}_{i+1} = W_i \vec{\varphi}_i \quad (i \in \mathbb{Z}), \quad (72)$$

where

$$U_i = A_i - B_i N_i^{-1}(\lambda + M_i), \quad W_i = -N_i^{-1}(\lambda + M_i) \quad (i \in \mathbb{Z}). \quad (73)$$

This type of 2×2 nonlinear chains is investigated in [1] in relation to Painlevé equations. It is not clear yet, however, how our system (72) can be related to the one employed there for obtaining P_{VI} .

In this paper, we have presented a new Lax pair for the sixth Painlevé equation in the framework of the loop algebra $\mathfrak{so}(8)[z, z^{-1}]$ of type $D_4^{(1)}$. We also explained how the affine Weyl group symmetry of P_{VI} can be obtained from the linear problem. We expect that the Lax pair discussed in this paper could be applied as well to other problems concerning Painlevé equations. Also, it would be an important problem to understand properly the relationship of our representation with various approaches to the sixth Painlevé equation as in [2].

References

- [1] V.E. Adler: Nonlinear chains and Painlevé equations, *Physica D* **73** (1994), 335–351.
- [2] R. Conte (Ed): *The Painlevé Property: One Century Later*, CRM Series in Mathematical Physics, Springer-Verlag, New York, 1999.
- [3] K. Iwasaki, K. Kimura, S. Shimomura and M. Yoshida: *From Gauss to Painlevé — A Modern Theory of Special Functions*, Aspects of Mathematics **E16**, Vieweg, 1991.
- [4] V.G. Kac: *Infinite dimensional Lie algebra*, Third Edition, Cambridge University Press, 1990.
- [5] M. Noumi and Y. Yamada: Affine Weyl groups, discrete dynamical systems and Painlevé equations, *Commun. Math. Phys.* **199**(1998), 281–295.
- [6] M. Noumi and Y. Yamada: Birational Weyl group action arising from a nilpotent Poisson algebra, in *Physics and Combinatorics 1999* (Eds. A.N. Kirillov, A. Tsuchiya and H. Umemura), pp. 287–319, World Scientific, 2001.
- [7] K. Okamoto: Study of the Painlevé equations I, *Ann. Math. Pura Appl.* **146**(1987), 337–381.